Math. 350-02 Spring, 2016 R. B. Kearfott

Final Exam Monday, May 2, 2016, 2:00PM-4:30PM

This exam is closed book, but you may use calculators. Make sure your name is on all pages. Show all work, and show it in a logical and organized manner. Each problem is worth 25 points.

1. Use the characteristic equation, as well as undetermined coefficients, to find the solution to the following initial value problem.

$$y'' + 2y' + 2y = 5\sin(t), \quad y(0) = -2, \ y'(0) = 1.$$

- 2. Consider $y(t) = 3\cos(2t) + 3\sqrt{3}\sin(2t)$.
 - (a) Rewrite y(t) in the form $y(t) = R \cos(\omega t \delta)$. (That is, find R, ω , and δ .)
 - (b) State the amplitude, natural frequency, and phase shift of y.
- 3. Write down the terms of the power series solution to the following initial value problem, up to and including the x^4 term.

$$y'' + 2y' + xy = x$$
, $y(0) = 1$, $y'(0) = -1$.

4. (Refer to Table 1 to do this problem.) Use Laplace transforms to find the solution to

$$y'' - y = 1$$
, $y(0) = 0$, $y'(0) = 0$.

$f(t) = \mathcal{L}^{-1}\{F(s)\}$	$F(s) = \mathcal{L}\{f(t)\}$	Notes
1 1 for the set 1 1 1 1	$\frac{1}{s}, \frac{s}{s} > 0$	
which $2. e^{at}$ is contained and a	$\frac{1}{s-a}, \qquad s > a$	Sec. 6.1; Ex. 5
3. t^n , $n = \text{positive integer}$	$\frac{n!}{s^{n+1}}, \qquad s>0$	Sec. 6.1; Prob. 27
$4. t^p, \qquad p > -1$	$\frac{\Gamma(p+1)}{s^{p+1}}, \qquad s > 0$	Sec. 6.1; Prob. 27
5. sin <i>at</i>	$\frac{a}{s^2 + a^2}, \qquad s > 0$	Sec. 6.1; Ex. 6
6. $\cos at$ and entropy of the constant of the	$\frac{s}{s^2 + a^2}, \qquad s > 0$	Sec. 6.1; Prob. 6
7. sinh <i>at</i>	$\frac{a}{s^2 - a^2}, \qquad s > a $	Sec. 6.1; Prob. 8
8. cosh <i>at</i>	$\frac{s}{s^2 - a^2}, \qquad s > a $	Sec. 6.1; Prob. 7
9. $e^{at} \sin bt$	$\frac{b}{(s-a)^2 + b^2}, \qquad s > a$	Sec. 6.1; Prob. 13
10. $e^{at}\cos bt$	$\frac{s-a}{(s-a)^2+b^2}, \qquad s>a$	Sec. 6.1; Prob. 14
11. $t^n e^{at}$, $n = \text{positive integer}$	$\frac{n!}{(s-a)^{n+1}}, \qquad s > a$	Sec. 6.1; Prob. 18
12. $u_c(t)$	$\frac{e^{-cs}}{s}, \qquad s > 0$	Sec. 6.3
13. $u_c(t)f(t-c)$	$e^{-cs}F(s)$	Sec. 6.3
14. $e^{ct}f(t)$	F(s-c)	Sec. 6.3
15. $f(ct)$	$\frac{1}{c}F\left(\frac{s}{c}\right), \qquad c > 0$	Sec. 6.3; Prob. 19
$16. \int_0^t f(t-\tau)g(\tau)d\tau$	F(s)G(s)	Sec. 6.6
17. $\delta(t-c)$	e^{-cs}	Sec. 6.5
18. $f^{(n)}(t)$	$s^{n}F(s) - s^{n-1}f(0) - \dots - f^{(n-1)}(0)$	Sec. 6.2
19. $(-t)^n f(t)$	$F^{(n)}(s)$	Sec. 6.2; Prob. 28

Table 1: Table of Laplace Transforms

319