Math. 350-02 Spring, 2018 R. B. Kearfott

Final Exam

Monday, April 30, 2018, 2:00PM to 4:30PM

This exam is closed book, but you may use calculators. Make sure your name is on all pages. Show all work, and show it in a logical and organized manner. (Do it carefully, and check it.) Each entire problem is worth 25 points.

- 1. A tank contains 400 gallons of water and 10 pounds of salt. Water containing a salt concentration of .25 pounds per gallon flows into the tank at a rate of 2 gallons per minute, and the mixture in the tank flows out at the same rate.
 - (a) Write down a differential equation relating the quantity Q of salt in the tank at time t and the derivative of Q.
 - (b) Solve that differential equation.
 - (c) Determine the amount of salt in the tank after a long time.
- 2. Consider the initial value problem

$$y'' + 12y' + 37y = \sin(t), \quad y(0) = 1, \ y'(0) = 0.$$

- (a) Solve the initial value problem. Be sure to carefully write down all steps in your solution process.
- (b) Write down the steady-state solution.
- 3. Write down the degree 5 polynomial approximation to the series solution to the following initial value problem.

$$y' + (x+1)y = 0, \quad y(0) = 1.$$

4. Use Laplace transforms to solve the following initial value problem. You may use the table on the back of this sheet.

$$y'' + 3y' + 2y = f(t), \quad y(0) = 1, \ y'(0) = 0, \quad \text{where} \quad f(t) = \left\{ \begin{array}{l} 0, \ 0 \le t < 1\\ 1, \ 1 \le t < 2\\ 0, \ t \ge 2 \end{array} \right\}.$$

TABLE 6.2.1 Elementary Laplace Transforms

$f(t) = \mathcal{L}^{-1}\{F(s)\}$	$F(s) = \mathcal{L}{f(t)}$	Notes
1 . 1 - $$	$\frac{1}{s}, \frac{1}{s > 0}$ is the anti-interval of the set	Sec. 6.1; Ex. 4
e^{at} . e^{at} . Solution we have 2 . e^{at} is following the Laplace transformed or 2 . 2	$\frac{1}{s-a}, \qquad s > a$	Sec. 6.1; Ex. 5
3. t^n , $n = \text{positive integer}$	$\frac{n!}{s^{n+1}}, \qquad s > 0$	Sec. 6.1; Prob. 27
4. t^p , $p > -1$	$\frac{\Gamma(p+1)}{s^{p+1}}, \qquad s > 0$	Sec. 6.1; Prob. 27
5. sin at	$\frac{a}{s^2 + a^2}, \qquad s > 0$	Sec. 6.1; Ex. 6
6. cos at anoi edimina	$\frac{s}{s^2 + a^2}, \qquad s > 0$	Sec. 6.1; Prob. 6
7. sinh <i>at</i>	$\frac{a}{s^2 - a^2}, \qquad s > a $	Sec. 6.1; Prob. 8
8. cosh <i>at</i>	$\frac{s}{s^2 - a^2}, \qquad s > a $	Sec. 6.1; Prob. 7
9. $e^{at} \sin bt$	$\frac{b}{(s-a)^2+b^2}, \qquad s>a$	Sec. 6.1; Prob. 13
10. $e^{at}\cos bt$	$\frac{s-a}{(s-a)^2+b^2}, \qquad s>a$	Sec. 6.1; Prob. 14
11. $t^n e^{at}$, $n = \text{positive integer}$	$\frac{n!}{(s-a)^{n+1}}, \qquad s > a$	Sec. 6.1; Prob. 18
12. $u_c(t)$	$\frac{e^{-cs}}{s}, \qquad s > 0$	Sec. 6.3
13. $u_c(t)f(t-c)$	$e^{-cs}F(s)$	Sec. 6.3
14. $e^{ct}f(t)$	F(s-c)	Sec. 6.3
15. $f(ct)$	$\frac{1}{c}F\left(\frac{s}{c}\right), \qquad c > 0$	Sec. 6.3; Prob. 19
$16. \int_0^t f(t-\tau)g(\tau)d\tau$	F(s)G(s)	Sec. 6.6
17. $\delta(t-c)$	e^{-cs}	Sec. 6.5
18. $f^{(n)}(t)$	$s^{n}F(s) - s^{n-1}f(0) - \dots - f^{(n-1)}(0)$	Sec. 6.2
$\frac{19. \ (-t)^n f(t)}{}$	$F^{(n)}(s)$	Sec. 6.2; Prob. 28